References

- Ana Carolina MelchiorsCassyano Januário Correr, Roberto Pontarolo, (2010) Felipe de Oliveira de Souza Santos, Rodrigo Augusto de Paula e Souza Arquivos Brasileiros de Cardiologia; 94(3):337-344)
- Albert, N. M (2008). Improving medication adherence in chronic cardiovascular disease. Journal of Critical Care Nurse28, No.3, 54-64.
- Balazovjech I Hnilica P Jr (1993), Compliance with antihypertensive treatment in consultation rooms for hypertensive patients. Journal of human hypertension7 (6):581–583
- Centers for Disease Control and Prevention (CDC) & National Institutes of Health (NIH). (2000). Healthy people 2010: 12 Heart disease and stroke. Retrieved August 14, 2010 from http://www.healthypeople.gov/Document/HTML/Volume1/12Heart.htm.
- Col, K. J. E. Fanale, et al (1990)"The role of medication compliance and adverse drug reactions in hospitalizations of the elderly." Arch Intern Med 150: 841-845.
- Dighore P N Patil PH et al (2009) Prescription patterns of antihypertensive drugs", Indian journal of hospital pharmacy 46 78-82.
- DiMatteo, M. R., P. J. Giordani, et al (2002) Patient adherence and medical treatment outcomes: a meta analysis." Med Care 40: 794-811.
- Dennehy, C. E., D. T. Kishi, et al (1996)"Drug-related illness in emergency departmentPatients." Am J Health-Syst Pharm 53: 1422-1426.
- Gray, S. LJ. E. Mahoney, et al. (1999) Adverse drug events in elderly patients recieving home health services following hospital discharge." Annals of pharmacotherapy 33: 1147-1153.
- Grueninger UJ (1995) Arterial hypertension: lessons from patient education. Patient Educ Couns. 26(1–3):37–55.
- Gupta R (1996), Jul al-Odat NA, Gupta VP. Hypertension epidemiology in India: meta-analysis of 50 year prevalence rates and blood pressure trends. J Hum Hypertens.;10(7):465-472.
- Gupta R, (2010) Guptha S. Strategies for initial management of hypertension. Indian J Med Res. Nov;132 5):531-542.

- Helper CD, Strand LM (1990) Opportunities and Responsibilities in pharmaceutical care. Am J Hosp Pharm; 47: 533- 543.
- Horne R (1993) One to be taken as directed: Reflections on non-adherence (non-compliance).Journal of Social and Administrative Pharmacy 10(4):150-156.
- Hansson LHedner T, Himmelman (1998) A. The growing importance of systolic blood pressure. Blood Press. 7:131–2.
- Irvine, J. B. Baker, et al (1999) Poor adherence to placebo or amiodarone therapy predicts mortality: results from the CAMIAT study." Psychosom Med 61: 566-575.
- Joint National Committee 7th report (JNC 2004) on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure NIH Publication No. 04-5230
- JNC VI (1997). The sixth report of the Joint National Committee on Prevention,Detection, Evaluation and Treatment of High Blood Pressure (JNC VI). Arch Intern Med. 1997;157:2413–46.
- Kaplan NM (2007). Systemic hypertension: therapy chap 41. In: Libby P, Bonow RO, Mann DL, Zipes DP, eds. Libby: Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 8th ed. Saunders.
- Kearney PMWhelton M, Reynolds K, Muntner P, Whelton PK, He J (2005) Global burden of hypertension: analysis of worldwide data. Lancet; 365: 217-23.
- Khosravi A R, Shirani SH, Shahrokhi SH, Mohammadifard N, Ansari R.(2006) Antihypertensive Drugs used by Hypertensive Patients In The Provincial Cities of Isfahan, Najafabad and Arak. ARYA Journal; 1(4): 275-80.
- Lawes, C (2008).. M. Hoorn, S. V., & Rodgers, A. Global burden of blood-pressurerelated disease, Lancet, 371(9623). 1513-15.
- Leupkar RV (1971). Patient adherence: A "risk factor" for cardiovascular disease.The Framington Study. JAMA; 215:1617-1625.
- Lima-Costa MF (2003), Veras R. Saúde Pública e Envelhecimento [Editorial]. Cad. Saúde Pública ;19(3):700–1.
- Moscal LLinfante AH (2005), Benjamin EJ, Berra K, Hayes SN, Walsch BW, et alNational study of physician awareness and adherence to cardiovascular disease prevention guidelines. Circulation;111:499-510

- Melinda Sue darling (2009) Non-pharmaceutical management of hypertension among middle aged and older adults. Master science thesis submitted to university of Arizona.www.Nursingarizona.edu/scripts/num=84141562351220991330
- Oscar A (2000), Carretero and Suzanne Oparil Essential Hypertension: Part I: Definition and EtiologyJ American Heart Association 101:329-335.
- Ong, K(2007). L., Cheung, B. M., Man, Y. B., Lau, C. P., & Lam, K. S.. Prevalence, awareness, treatment, and control of hypertension among United States adults1999-2004. Hypertension,49(1). 69-75.
- Osterberg L(2005), Blaschke T. Adherence to medication. N Engl J Med Aug;353(5):487-497.
- RichardE. Klabunde, Cardiovascular Pharmacology Concepts http://www.cvpharmacology.com/
- Rodgers A (2000), Lawes C, MacMahon S. Reducing the global burden of blood pressure related cardiovascular disease. J Hypertens; 18:S3-6.
- Roccella EJ (1988), Horan MJ. The National High Blood Pressure Education Program: measuring progress and assessing its impact. Health Psychol.;(suppl):297–303.
- Sewitch MJ (2004), Dobkin PL, Bernatsky S, Baron M, Starr M, Cohen M, et al. Medication non-adherence in women with fibromyalgia. Rheumatology (Oxford) May;43 (5):648-654.
- Sackett DL (1979), Snow JC. The magnitude of compliance and non compliance. IN Haynes NRB, Taylor DW, Sackett DL, eds. Compliance in Health Care. Baltimore: John Hopkins University Press;:11-22.
- Schlenk, E. A (2001), Burke, L.E., & Rand, C. Behavioral strategies to improve medication-taking compliance. In Burke, L.E., Ockene, I.S., eds. Compliance in Healthcare and Research;Armonk, NY: Futura Publishing Company. 57- 70.
- Svetkey LP (2005) et al. Effect of life style modification on BP by Race, Sex, Hypertension status and Age. Journal of human hypertension. January; 1: 21-31.
- Saounatsu M (2001), Patsi O, Fasoi G, Stylianou M, Kavga A, Economou O et al. The influence of the hypertensive patient's education in compliance with their medication Public Health Nursing; 18(6):436-442.

- Vasan RS, (2001)Larson MG, Leip EP, Kannel WB, Levy DAssessment of frequency of progression to hypertension in non-hypertensive participants in the Framingham Heart Study: a cohort study Lancet 17;358(9294):1682-6.
- VK Agrawal (2008), R Bhalwar, Basannar prevalence and determinants of hypertension in a rural community MJAFI, Vol. 64, No. 1, 21-25
- Viera AJ (2008), Cohen LW, Mitchell CM, Sloane PD. High blood pressure knowledge among primary care patients with known hypertension: a North Carolina Family Medicine Research Network (NC-FM-RN) study. J Am Board Fam Med; 21(4): 300-8.
- WHO (1996), Hypertension Control: Report of a WHO expert committee. WHO Technical Report Series No 862, World Health Organization, Geneva,
- World Health Organization, International Society of Hypertension Writing Group. (2003). World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. Journal of Hypertension, 21(11). 1983-1992.
- WHO (1996), Hypertension Control: Report of a WHO expert committee. WHO Technical Report Series No 862, World Health Organization, Geneva.
- White, T. J. (1999), A. Arakelian, et al. "Counting the costs of drug-related adverse events." Pharmacoeconomics 15: 445-458.
- Williams B (2004), Poulter NR, Brown MJ, Davis M, McInnes GT, Potter JF et al. British hypertension society guidelines for hypertension management 2004 (BHS- IV). BMJ; 328:634–40.